

Show *all* of your work and *explain* your answers fully. There is a total of 100 possible points. Partial credit is proportional to the quality of your explanation. You may use Sage to manipulate and row-reduce matrices. No other use of Sage may be used as justification for your answers. When you use Sage be sure to explain your input and show any relevant output (rather than just describing salient features).

1.  $P_2$  is the vector space of polynomials with degree at most 2. Find a basis for the subspace  $W$ . (You may assume that  $W$  is a subspace.) (15 points)

$$W = \{a + bx + cx^2 \mid a + 3b = 0, 2a + b - 5c = 0\} \subseteq P_2$$

2. The set  $B = \{(2, 1), (3, 4)\}$  is a basis for the crazy vector space,  $C$ . (You may assume this.) Express the vector  $(4, 0)$  as a linear combination of the basis vectors. (The operations of  $C$  are reproduced below.) (15 points)

$$(x_1, x_2) + (y_1, y_2) = (x_1 + y_1 + 1, x_2 + y_2 + 1) \quad \alpha(x_1, x_2) = (\alpha x_1 + \alpha - 1, \alpha x_2 + \alpha - 1)$$



3. The following are subsets of the vector space of  $2 \times 2$  matrices,  $M_{22}$ . Decide which are bases for  $M_{22}$ . Give complete justification for your answers, since a simple yes/no answer will get no credit. (25 points)

(a)  $J = \left\{ \begin{bmatrix} 3 & 6 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} -5 & 3 \\ -4 & 1 \end{bmatrix}, \begin{bmatrix} -3 & 0 \\ 2 & 6 \end{bmatrix}, \begin{bmatrix} 2 & 2 \\ -8 & 4 \end{bmatrix}, \begin{bmatrix} 7 & 3 \\ 5 & -3 \end{bmatrix} \right\}$

(b)  $K = \left\{ \begin{bmatrix} 1 & -2 \\ 3 & -7 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 2 & -3 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 3 & -6 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 2 & -2 \end{bmatrix} \right\}$

(c)  $L = \left\{ \begin{bmatrix} 1 & 0 \\ -3 & -7 \end{bmatrix}, \begin{bmatrix} 2 & 1 \\ -2 & -8 \end{bmatrix}, \begin{bmatrix} -1 & -1 \\ 0 & 3 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & -2 \end{bmatrix} \right\}$

4. Suppose that  $A$  is a  $5 \times 7$  matrix, which is row-equivalent to a matrix in reduced row-echelon form with 3 pivot columns. (15 points)

(a) Compute the rank of  $A$ .

(b) Compute the nullity of  $A$ .

(c) Compute the rank of  $A^t$ .

(d) Compute the nullity of  $A^t$ .



5. Demonstrate the use of the three parts of Theorem TSS to show that the set  $U$  is a subspace of the vector space  $\mathbb{C}^2$ . (15 points)

$$U = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} \mid 2a + 5b = 0 \right\} \subseteq \mathbb{C}^2$$

6. Suppose that  $V$  is a vector space,  $\mathbf{v} \in V$ , and  $\alpha \in \mathbb{C}$ . Prove that if  $\alpha\mathbf{v} = \mathbf{0}$ , then  $\alpha = 0$  or  $\mathbf{v} = \mathbf{0}$ . (Your proof should be more than simply quoting a result from the book.) (15 points)

