

Show *all* of your work and *explain* your answers fully. There is a total of 100 possible points.

Partial credit is proportional to the quality of your explanation. You may use Sage to row-reduce matrices, compute determinants, and compute eigenvalues, eigenmatrices and eigenspaces. No other use of Sage may be used as justification for your answers. When you use Sage be sure to explain your input and show any relevant output (rather than just describing salient features).

1. Let \mathbf{u} be an element of the vector space of 2×2 symmetric matrices, S_{22} , and let B be a basis for the vector space. Compute the vector representation of \mathbf{u} relative to B , $\rho_B(\mathbf{u})$. (15 points)

$$\mathbf{u} = \begin{bmatrix} 2 & 4 \\ 4 & -3 \end{bmatrix} \quad B = \left\{ \begin{bmatrix} 1 & -2 \\ -2 & 0 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} 6 & -8 \\ -8 & 5 \end{bmatrix} \right\}$$

2. E and F are bases for the vector space \mathbb{C}^2 . Compute the change-of-basis matrix, $C_{E,F}$. (15 points)

$$E = \left\{ \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \end{bmatrix} \right\} \quad F = \left\{ \begin{bmatrix} -1 \\ 11 \end{bmatrix}, \begin{bmatrix} 1 \\ -18 \end{bmatrix} \right\}$$



3. Consider the linear transformation T from the vector space of 1×2 matrices, M_{12} , to the vector space of polynomials with largest degree 1, P_1 . B and E are bases of M_{12} , C and F are bases of P_1 . (20 points)

$$T: M_{12} \rightarrow P_1, \quad T\left(\begin{bmatrix} a & b \end{bmatrix}\right) = (2a + 3b) + (a + b)x$$

$$B = \left\{ \begin{bmatrix} 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \end{bmatrix} \right\} \quad C = \{1, x\}$$

$$E = \left\{ \begin{bmatrix} 1 & 1 \end{bmatrix}, \begin{bmatrix} 3 & 4 \end{bmatrix} \right\} \quad F = \{2 + x, 3 + x\}$$

$$\mathbf{u} = \begin{bmatrix} 3 & 1 \end{bmatrix}$$

- (a) Compute $T(\mathbf{u})$ using the definition above.

- (b) Compute the matrix representation of T relative to B and C .

- (c) Compute $T(\mathbf{u})$, using the matrix representation from (b) and two vector representation linear transformations (ρ_X).

- (d) Compute the matrix representation of T relative to E and F .

- (e) Compute $T(\mathbf{u})$, using the matrix representation from (d) and two vector representation linear transformations (ρ_X).



4. Consider the invertible linear transformation T from the vector space \mathbb{C}^3 to the vector space of polynomials P_2 . Compute an explicit formula for the inverse of T , the linear transformation $T^{-1}: P_2 \rightarrow \mathbb{C}^3$. Do this using techniques from Chapter R, not with techniques from Chapter LT, such as computing pre-images. Do not skip any steps for dealing with \mathbb{C}^3 that might seem trivial, they must be described explicitly. (15 points)

$$T: \mathbb{C}^3 \rightarrow P_2, \quad T \left(\begin{bmatrix} a \\ b \\ c \end{bmatrix} \right) = (-5a - 2b + 2c) + (2a + b - c)x + (-3a - b)x^2$$

5. Find a basis for P_2 , the vector space of polynomials of degree at most 2, which will yield a diagonal matrix representation for the linear transformation R . (15 points)

$$R: P_2 \rightarrow P_2, \quad R(a + bx + cx^2) = (-5a + 4b) + (-8a + 7b)x + (4a - 4b - c)x^2$$

